

AD-A133 336

ON BAYES AND EMPIRICAL BAYES RULES FOR SELECTING GOOD  
POPULATIONS(U) PURDUE UNIV LAFAYETTE IN DEPT OF  
STATISTICS S S GUPTA ET AL. SEP 83 TR-83-37


1/1

UNCLASSIFIED


N00014-75-C-0455

F/G 12/1

NL



END  
DATE  
FILMED  
10 83  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A133 336

(12)

# PURDUE UNIVERSITY



DEPARTMENT OF STATISTICS

OCT 1 1 1983

A

DTIC FILE COPY

For information only  
Do not use for identification  
of statistical data.

83 10 06 001

**ON BAYES AND EMPIRICAL BAYES RULES  
FOR SELECTING GOOD POPULATIONS\***


by

**Shanti S. Gupta and Lii-Yuh Leu  
Purdue University**

**Technical Report #83-37**

**Department of Statistics  
Purdue University**

**September 1983**



This document has been approved  
for public release and sale; its  
distribution is unlimited.

**\*This research was supported by the Office of Naval Research Contract  
N00014-75-C-0455 at Purdue University. Reproduction in whole or in part  
is permitted for any purpose of the United States Government.**

-a-

ON BAYES AND EMPIRICAL BAYES RULES  
FOR SELECTING GOOD POPULATIONS

by

Shanti S. Gupta and Lii-Yuh Leu  
Purdue University

Abstract

This paper deals with the problem of selecting all populations which are close to a control or standard. A general Bayes rule for the above problem is derived. Empirical Bayes rules are derived when the populations are assumed to be uniformly distributed. Under some conditions on the marginal and prior distributions, the rate of convergence of the empirical Bayes risk to the minimum Bayes risk is investigated. The rate of convergence is shown to be  $n^{-\delta/3}$  for some  $\delta$ ,  $0 < \delta < 2$ .

Key words: Bayes rules, empirical Bayes rules, selection procedures, asymptotically optimal, rate of convergence.

\*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

-2-

ON BAYES AND EMPIRICAL BAYES RULES  
FOR SELECTING GOOD POPULATIONS

by

Shanti S. Gupta and Lii-Yuh Leu  
Purdue University

Abstract

This paper deals with the problem of selecting all populations which are close to a control or standard. A general Bayes rule for the above problem is derived. Empirical Bayes rules are derived when the populations are assumed to be uniformly distributed. Under some conditions on the marginal and prior distributions, the rate of convergence of the empirical Bayes risk to the minimum Bayes risk is investigated. The rate of convergence is shown to be  $n^{-\delta/3}$  for some  $\delta$ ,  $0 < \delta < 2$ .

Key words: Bayes rules, empirical Bayes rules, selection procedures, asymptotically optimal, rate of convergence.

\*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ON BAYES AND EMPIRICAL BAYES RULES  
FOR SELECTING GOOD POPULATIONS

by

Shanti S. Gupta and Lii-Yuh Leu  
Purdue University



Decision For	
1. H&I	<input checked="" type="checkbox"/>
2. H&I	<input type="checkbox"/>
3. H&I	<input type="checkbox"/>
4. H&I	<input type="checkbox"/>
5. H&I	<input type="checkbox"/>
6. H&I	<input type="checkbox"/>
7. H&I	<input type="checkbox"/>
8. H&I	<input type="checkbox"/>
9. H&I	<input type="checkbox"/>
10. H&I	<input type="checkbox"/>
11. H&I	<input type="checkbox"/>
12. H&I	<input type="checkbox"/>
13. H&I	<input type="checkbox"/>
14. H&I	<input type="checkbox"/>
15. H&I	<input type="checkbox"/>
16. H&I	<input type="checkbox"/>
17. H&I	<input type="checkbox"/>
18. H&I	<input type="checkbox"/>
19. H&I	<input type="checkbox"/>
20. H&I	<input type="checkbox"/>
21. H&I	<input type="checkbox"/>
22. H&I	<input type="checkbox"/>
23. H&I	<input type="checkbox"/>
24. H&I	<input type="checkbox"/>
25. H&I	<input type="checkbox"/>
26. H&I	<input type="checkbox"/>
27. H&I	<input type="checkbox"/>
28. H&I	<input type="checkbox"/>
29. H&I	<input type="checkbox"/>
30. H&I	<input type="checkbox"/>
31. H&I	<input type="checkbox"/>
32. H&I	<input type="checkbox"/>
33. H&I	<input type="checkbox"/>
34. H&I	<input type="checkbox"/>
35. H&I	<input type="checkbox"/>
36. H&I	<input type="checkbox"/>
37. H&I	<input type="checkbox"/>
38. H&I	<input type="checkbox"/>
39. H&I	<input type="checkbox"/>
40. H&I	<input type="checkbox"/>
41. H&I	<input type="checkbox"/>
42. H&I	<input type="checkbox"/>
43. H&I	<input type="checkbox"/>
44. H&I	<input type="checkbox"/>
45. H&I	<input type="checkbox"/>
46. H&I	<input type="checkbox"/>
47. H&I	<input type="checkbox"/>
48. H&I	<input type="checkbox"/>
49. H&I	<input type="checkbox"/>
50. H&I	<input type="checkbox"/>
51. H&I	<input type="checkbox"/>
52. H&I	<input type="checkbox"/>
53. H&I	<input type="checkbox"/>
54. H&I	<input type="checkbox"/>
55. H&I	<input type="checkbox"/>
56. H&I	<input type="checkbox"/>
57. H&I	<input type="checkbox"/>
58. H&I	<input type="checkbox"/>
59. H&I	<input type="checkbox"/>
60. H&I	<input type="checkbox"/>
61. H&I	<input type="checkbox"/>
62. H&I	<input type="checkbox"/>
63. H&I	<input type="checkbox"/>
64. H&I	<input type="checkbox"/>
65. H&I	<input type="checkbox"/>
66. H&I	<input type="checkbox"/>
67. H&I	<input type="checkbox"/>
68. H&I	<input type="checkbox"/>
69. H&I	<input type="checkbox"/>
70. H&I	<input type="checkbox"/>
71. H&I	<input type="checkbox"/>
72. H&I	<input type="checkbox"/>
73. H&I	<input type="checkbox"/>
74. H&I	<input type="checkbox"/>
75. H&I	<input type="checkbox"/>
76. H&I	<input type="checkbox"/>
77. H&I	<input type="checkbox"/>
78. H&I	<input type="checkbox"/>
79. H&I	<input type="checkbox"/>
80. H&I	<input type="checkbox"/>
81. H&I	<input type="checkbox"/>
82. H&I	<input type="checkbox"/>
83. H&I	<input type="checkbox"/>
84. H&I	<input type="checkbox"/>
85. H&I	<input type="checkbox"/>
86. H&I	<input type="checkbox"/>
87. H&I	<input type="checkbox"/>
88. H&I	<input type="checkbox"/>
89. H&I	<input type="checkbox"/>
90. H&I	<input type="checkbox"/>
91. H&I	<input type="checkbox"/>
92. H&I	<input type="checkbox"/>
93. H&I	<input type="checkbox"/>
94. H&I	<input type="checkbox"/>
95. H&I	<input type="checkbox"/>
96. H&I	<input type="checkbox"/>
97. H&I	<input type="checkbox"/>
98. H&I	<input type="checkbox"/>
99. H&I	<input type="checkbox"/>
100. H&I	<input type="checkbox"/>

1. Introduction

Empirical Bayes rules have been considered for multiple decision problems by Deely (1965), Van Ryzin (1970), Van Ryzin and Susarla (1977), Singh (1977), and Gupta and Hsiao (1981). Most of the papers are concerned with the selection of the best population where best is usually defined in terms of the largest or smallest unknown parameter. Gupta and Hsiao (1981) considered the problem which is concerned with the selection of populations better than a control. In some practical applications, one may be interested in selecting populations which are close to a control. We will consider this kind of problem in this paper.

In Section 2, we propose a general Bayes rule for selecting good populations. In Section 3, assuming that the populations are uniformly distributed, empirical Bayes rules are derived for both the known control parameter and the unknown control parameter cases. Under some conditions on the marginal and prior distributions, the rate of convergence of the empirical Bayes risk to the minimum Bayes risk is investigated. The rate of convergence is shown to be  $n^{-\delta/3}$  for some  $\delta$ ,  $0 < \delta < 2$ .

---

\*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

## 2. A General Bayes Rule for Selecting Good Populations

Let  $\pi_0, \pi_1, \dots, \pi_k$  be  $(k+1)$  independent populations which are characterized by parameters  $\theta_0, \theta_1, \dots, \theta_k$ , respectively. Assume that  $\pi_0$  is the control population with parameter  $\theta_0$  which may be known or unknown. When  $\theta_0$  is unknown, let  $\underline{\theta} = (\theta_0, \theta_1, \dots, \theta_k)$  and  $\underline{X} = (X_0, X_1, \dots, X_k)$  where  $X_i$  is an observation from  $\pi_i$ ,  $i = 0, 1, \dots, k$ . When  $\theta_0$  is known, no observation from population  $\pi_0$  is taken, and  $\theta_0, X_0$  are deleted from  $\underline{\theta}$  and  $\underline{X}$ , respectively. When there is no confusion,  $\underline{\theta}$  and  $\underline{X}$  are used to represent either case. We define population  $\pi_i$  to be a good population if  $|\theta_i - \theta_0| < \Delta$  and a bad population if  $|\theta_i - \theta_0| \geq \Delta$ , where  $\Delta > 0$  is a pre-assigned constant. Our goal is to find a Bayes rule which selects all good populations and rejects bad ones. We assume that given  $\theta_i$ ,  $X_i$  has probability density function  $f(x_i | \theta_i)$  with respect to a  $\sigma$ -finite measure  $\mu$ , for  $i = 0, 1, \dots, k$ , and  $\underline{\theta}$  has a prior distribution

$G(\underline{\theta}) = \prod_{i=0}^k G_i(\theta_i)$  on the parameter space  $\Omega$ . Let  $G = \{s | s \subseteq \{1, 2, \dots, k\}\}$  be the action space and let

$$(2.1) \quad L(\underline{\theta}, s) = \sum_{i \in s} \{c_1(\theta_0 - \Delta - \theta_i) I_{\{\theta_i \leq \theta_0 - \Delta\}}(\theta_i) + \\ c_2(\theta_i - \theta_0 - \Delta) I_{\{\theta_0 + \Delta \leq \theta_i\}}(\theta_i)\} + \sum_{i \notin s} \{c_3(\theta_i - \theta_0 + \Delta) I_{\{\theta_0 - \Delta < \theta_i \leq \theta_0\}}(\theta_i) \\ + c_4(\theta_0 + \Delta - \theta_i) I_{\{\theta_0 < \theta_i < \theta_0 + \Delta\}}(\theta_i)\}$$

be the loss function defined on  $\Omega \times G$ , where  $c_i$ ,  $i = 1, 2, 3, 4$  are positive constants and  $I$  is the indicator function. The Bayes risk with respect to  $G$  can be expressed as

$$(2.2) \quad r(G, s) = \int_{\mathcal{X}} \int_{\Omega} L(\underline{\theta}, s) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) d\mu(\underline{x}),$$

where  $\mathcal{X}$  is the sample space and  $f(\underline{x} | \underline{\theta}) = \prod_i f(x_i | \theta_i)$ .



Since the action space is finite, attention can be restricted to the non-randomized rules for deriving the Bayes rules. For a non-randomized decision function  $\delta: \mathcal{X} \rightarrow G$ , the corresponding Bayes risk with respect to  $G$  is given by

$$(2.3) \quad r(G, \delta) = \int_{\mathcal{X}} \int_{\Omega} L(\underline{\theta}, \delta(\underline{x})) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) d\mu(\underline{x}).$$

In the sequel we consider the special case where  $c_1 = c_2 = c_3 = c_4 =$  constant which can be taken to be unity without loss of generality. If  $\phi$  is the empty set, we have

$$L(\underline{\theta}, \phi) = \sum_{i=1}^k \{ (\theta_i - \theta_0 + \Delta) I_{\{\theta_0 - \Delta < \theta_i \leq \theta_0\}}(\theta_i) + (\theta_0 + \Delta - \theta_i) I_{\{\theta_0 < \theta_i < \theta_0 + \Delta\}}(\theta_i) \},$$

and (2.1) can be expressed as

$$(2.4) \quad L(\underline{\theta}, s) = L(\underline{\theta}, \phi) + \sum_{i \in s} \{ (\theta_0 - \Delta - \theta_i) I_{\{\theta_i \leq \theta_0\}}(\theta_i) + (\theta_i - \theta_0 - \Delta) I_{\{\theta_0 < \theta_i\}}(\theta_i) \}.$$

Hence, for any  $\delta$ , we have

$$(2.5) \quad \begin{aligned} r(G, \delta) - r(G, \phi) &= \int_{\mathcal{X}} \sum_{i \in \delta(\underline{x})} \left\{ \int_{\Omega} (\theta_0 - \Delta - \theta_i) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) + \right. \\ &\quad \left. 2 \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) d\mu(\underline{x}) \right\}. \end{aligned}$$

From (2.5),  $\delta_B(\underline{x})$  is given by  $i \in \delta_B(\underline{x})$  if

$$(2.6) \quad \int_{\Omega} (\theta_0 - \Delta - \theta_i) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) + 2 \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) f(\underline{x} | \underline{\theta}) dG(\underline{\theta}) < 0,$$

then  $\delta_B(\underline{x})$  is a Bayes rule with respect to  $G$ .

Let  $m_i(x_i) = \int_{\Omega} f(x_i|\theta_i) dG_i(\theta_i)$  be the marginal distribution of  $X_i$ ,  $\pi(\theta_i|x_i)$  be the posterior distribution of  $\theta_i$  given  $X_i = x_i$ , and  $E(\theta_i|x_i)$  be the expected value of  $\theta_i$  given  $X_i = x_i$ . If  $m_i(x_i) > 0$  for all  $x_i$ , then (2.6) is equivalent to

$$(2.7) \quad (\theta_0 - \Delta) - E(\theta_i|x_i) + 2 \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) \pi(\theta_i|x_i) d\theta_i < 0$$

if  $\theta_0$  is known, or

$$(2.8) \quad E(\theta_0|x_0) - E(\theta_i|x_i) - \Delta + 2 \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) \pi(\theta_i|x_i) \pi(\theta_0|x_0) d\theta_i d\theta_0 < 0$$

if  $\theta_0$  is unknown.

From the above discussion, we have the following main result:

Theorem 2.1. Under the loss function (2.4), the Bayes rule  $\delta_B(\underline{x})$  with respect to  $G$  is given by

- (a) If  $\theta_0$  is known, then  $i \in \delta_B(\underline{x})$  if the inequality (2.7) holds.
- (b) If  $\theta_0$  is unknown, then  $i \in \delta_B(\underline{x})$  if the inequality (2.8) holds

#### An Application:

Suppose that

$$(2.9) \quad f(x_i|\theta_i) = e^{-\theta_i} \theta_i^{x_i} / (x_i!), \quad x_i = 0, 1, \dots, \theta_i > 0$$

and  $\theta_i$  has a prior distribution  $g_i(\theta_i) = G_i'(\theta_i)$  which is given by

$$(2.10) \quad g_i(\theta_i) = \frac{\alpha_i^{\alpha_i} \theta_i^{\alpha_i-1}}{\beta_i^{\alpha_i}} e^{-\beta_i \theta_i} / \Gamma(\alpha_i) I_{(0,\infty)}(\theta_i),$$

where  $\alpha_i > 0$  and  $\beta_i > 0$  are known. Then

$$(2.11) \quad \pi(\theta_i|x_i) = \frac{(1+\beta_i)^{x_i+\alpha_i} \theta_i^{x_i+\alpha_i-1} e^{-\theta_i(1+\beta_i)}}{\Gamma(x_i+\alpha_i)}$$

and

$$(2.12) \quad E(\theta_i|x_i) = (x_i + \alpha_i) / (1 + \beta_i).$$

Lemma 2.2. If  $\pi(\theta_i | x_i)$  is defined by (2.11), then

$$\begin{aligned} & \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) \pi(\theta_i | x_i) d\theta_i \\ &= \frac{x_i + \alpha_i}{1 + \beta_i} \{1 - \Gamma(\theta_0(1 + \beta_i); x_i + \alpha_i + 1)\} - \theta_0 \{1 - \Gamma(\theta_0(1 + \beta_i); x_i + \alpha_i)\}, \end{aligned}$$

where  $\theta_0 > 0$  is known and

$$\Gamma(a; \alpha) = \int_0^a \frac{x^{\alpha-1}}{\Gamma(\alpha)} e^{-x} dx, \quad a > 0, \quad \alpha > 0.$$

Proof. Proof is simple and hence omitted.

Lemma 2.3. If  $\pi(\theta_i | x_i)$  is defined by (2.11) and  $\beta_i = \beta$ ,  $i = 0, 1, \dots, k$  and  $\theta_0$  is unknown, then

$$\begin{aligned} & \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) \pi(\theta_i | x_i) \pi(\theta_0 | x_0) d\theta_i d\theta_0 \\ &= \frac{(x_i + \alpha_i + x_0 + \alpha_0)}{1 + \beta} I\left(\frac{1}{2}; x_0 + \alpha_0, x_i + \alpha_i\right) - \frac{2(x_0 + \alpha_0)}{1 + \beta} I\left(\frac{1}{2}; x_0 + \alpha_0 + 1, x_i + \alpha_i\right), \end{aligned}$$

where

$$I(z; \alpha, \beta) = \int_0^z \frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1} dx, \quad \alpha > 0, \quad \beta > 0,$$

and

$$B(\alpha, \beta) = \Gamma(\alpha)\Gamma(\beta)/\Gamma(\alpha+\beta).$$

Proof.  $\int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) \pi(\theta_i | x_i) \pi(\theta_0 | x_0) d\theta_i d\theta_0$

$$\begin{aligned} &= \int_0^\infty \int_{\theta_0}^\infty (\theta_i - \theta_0) \frac{(1+\beta)^{x_i + \alpha_i + x_0 + \alpha_0}}{\Gamma(x_i + \alpha_i) \Gamma(x_0 + \alpha_0)} \theta_i^{x_i + \alpha_i - 1} \theta_0^{x_0 + \alpha_0 - 1} e^{-(\theta_i + \theta_0)(1+\beta)} d\theta_i d\theta_0 \\ &= \int_0^\infty \int_1^\infty \frac{(1+\beta)^{x_i + \alpha_i + x_0 + \alpha_0}}{\Gamma(x_i + \alpha_i) \Gamma(x_0 + \alpha_0)} \theta_0^{x_i + \alpha_i + x_0 + \alpha_0} (u-1) u^{x_i + \alpha_i - 1} e^{-(1+u)\theta_0(1+\beta)} du d\theta_0 \end{aligned}$$

$$\begin{aligned}
&= \int_0^{\infty} \frac{(x_i + \alpha_i + x_0 + \alpha_0)}{(1+\beta)B(x_0 + \alpha_0, x_i + \alpha_i)} (u-1)u^{x_i + \alpha_i - 1} (1+u)^{-(x_i + \alpha_i + x_0 + \alpha_0 + 1)} du \\
&= \frac{(x_i + \alpha_i + x_0 + \alpha_0)}{(1+\beta)B(x_0 + \alpha_0, x_i + \alpha_i)} \int_0^1 (1-v)(v+1)^{-(x_i + \alpha_i + x_0 + \alpha_0 + 1)} v^{x_0 + \alpha_0 - 1} dv \\
&= \frac{(x_i + \alpha_i + x_0 + \alpha_0)}{(1+\beta)B(x_0 + \alpha_0, x_i + \alpha_i)} \int_0^{\frac{1}{2}} (1-2s)(1-s)^{x_i + \alpha_i - 1} s^{x_0 + \alpha_0 - 1} ds \\
&= \frac{(x_i + \alpha_i + x_0 + \alpha_0)}{1+\beta} I\left(\frac{1}{2}; x_0 + \alpha_0, x_i + \alpha_i\right) - \frac{2(x_0 + \alpha_0)}{1+\beta} I\left(\frac{1}{2}; x_0 + \alpha_0 + 1, x_i + \alpha_i\right).
\end{aligned}$$

From Theorem 2.1, Lemma 2.2 and Lemma 2.3, we have the following theorem:

**Theorem 2.4.** If  $f(x_i | \theta_i)$  is defined by (2.9) and  $g_i(\theta_i)$  is defined by (2.10). Under the loss function (2.4), the Bayes rule  $\delta_B(\underline{x})$  is given by

(a) If  $\theta_0$  is known, then  $i \in \delta_B(\underline{x})$  if

$$\frac{x_i + \alpha_i}{1 + \beta_i} \{1 - 2\Gamma(\theta_0(1 + \beta_i); x_i + \alpha_i + 1)\} - \theta_0 \{1 - 2\Gamma(\theta_0(1 + \beta_i); x_i + \alpha_i)\} < \Delta.$$

(b) If  $\theta_0$  is unknown and  $\beta_i = \beta$ ,  $i = 0, 1, \dots, k$ , then  $i \in \delta_B(\underline{x})$  if

$$\begin{aligned}
&\frac{x_i + \alpha_i}{1 + \beta} \{2I(\frac{1}{2}; x_0 + \alpha_0, x_i + \alpha_i) - 1\} + \frac{x_0 + \alpha_0}{1 + \beta} \{1 + \\
&2I(\frac{1}{2}; x_0 + \alpha_0, x_i + \alpha_i) - 4I(\frac{1}{2}; x_0 + \alpha_0 + 1, x_i + \alpha_i)\} < \Delta.
\end{aligned}$$

### 3. Empirical Bayes Rules for Uniform Populations

In this section we will assume that  $X_i$  has probability density function

$f(x_i | \theta_i) = \frac{1}{\theta_i} I_{(0, \theta_i)}(x_i)$ , where  $\theta_i > 0$  is unknown. Suppose that  $\underline{\theta}$  has a prior distribution  $G(\underline{\theta}) = \pi G_i(\theta_i)$  on  $\Omega$  and  $G_i$  has a continuous probability

density function  $g_i$  and  $g_i$  is positive. Let  $m_i(x_i)$ ,  $M_i(x_i)$  be the marginal pdf and cdf of  $X_i$ , respectively. Then

$$(3.1) \quad m_i(x_i) = \int_{x_i}^{\infty} \frac{1}{\theta_i} dG_i(\theta_i)$$

and

$$(3.2) \quad M_i(x_i) = x_i m_i(x_i) + G_i(x_i).$$

From (3.2), we have

$$(3.3) \quad G_i(x_i) = M_i(x_i) - x_i m_i(x_i).$$

It follows that

$$(3.4) \quad \int_a^b \frac{1}{\theta_i} dG_i(\theta_i) = m_i(a) - m_i(b)$$

and

$$\int_a^{\infty} \frac{1}{\theta_i} dG_i(\theta_i) = m_i(a)$$

for any  $x_i \leq a < b < \infty$ .

### 3.1. $\theta_0$ known

In the case where  $\theta_0$  is known, let

$$(3.5) \quad \Delta_{G_i}(x_i) = (\theta_0 - \Delta) m_i(x_i) - \int_{x_i}^{\infty} dG_i(\theta_i) + 2 \int_{(x_i, \infty) \cap (\theta_0, \infty)} (\theta_i - \theta_0) f(x_i | \theta_i) dG_i(\theta_i).$$

From (2.6), we have  $i \in \delta_B(\underline{x})$  if  $\Delta_{G_i}(x_i) < 0$ . If  $x_i \leq \theta_0$ ,

$$\begin{aligned} (3.6) \quad \Delta_{G_i}(x_i) &= (\theta_0 - \Delta) m_i(x_i) - \int_{x_i}^{\infty} dG_i(\theta_i) + 2 \int_{\theta_0}^{\infty} (\theta_i - \theta_0) \frac{1}{\theta_i} dG_i(\theta_i) \\ &= (\theta_0 - \Delta - x_i) m_i(x_i) + 1 - 2M_i(\theta_0) + M_i(x_i) \\ &= \Delta_{1, G_i}(x_i) \quad (\text{say}). \end{aligned}$$

If  $x_i > \theta_0$ ,

$$\begin{aligned}
(3.7) \quad \Delta_{G_i}(x_i) &= (\theta_0 - \Delta)m_i(x_i) - \int_{x_i}^{\infty} dG_i(\theta_i) + 2 \int_{x_i}^{\infty} (\theta_i - \theta_0) \frac{1}{\theta_i} dG_i(\theta_i) \\
&= (x_i - \theta_0 - \Delta)m_i(x_i) + 1 - M_i(x_i) \\
&= \Delta_{2,G_i}(x_i) \quad (\text{say}).
\end{aligned}$$

Therefore

$$(3.8) \quad \delta_B(\underline{x}) = \{i | x_i \leq \theta_0, \Delta_{1,G_i}(x_i) < 0\} \cup \{i | x_i > \theta_0, \Delta_{2,G_i}(x_i) < 0\}.$$

Remarks:

(1)  $\Delta_{G_i}(x_i)$  is strictly decreasing for  $0 < x_i < \theta_0 - \Delta$ , strictly increasing for  $\theta_0 - \Delta < x_i < \theta_0 + \Delta$ , and strictly decreasing for  $\theta_0 + \Delta < x_i$  (we assume that  $\theta_0 - \Delta > 0$ ).

(2) If  $x_i \geq \theta_0 + \Delta$ , then  $\Delta_{G_i}(x_i) \geq 0$ . Hence  $i \notin \delta_B(\underline{x})$  if  $x_i \geq \theta_0 + \Delta$ .

(3) If  $G_i$  is such that  $1 - 2M_i(\theta_0) + M_i(\theta_0 - \Delta) \geq 0$ , then  $\delta_B(\underline{x}) = \phi$ .

Otherwise,  $i \in \delta_B(\underline{x})$  if  $(\theta_0 - \Delta) - d_1 < x_i < (\theta_0 + \Delta) - d_2$ , for some positive real numbers  $d_1$  and  $d_2$ . Hence this type of selection rules are Bayes rules relative to some prior distribution.

If  $G$  is unknown, the Bayes rules are not obtainable. In this case, we consider a sequence  $(\underline{x}_1, \underline{\Delta}_1), (\underline{x}_2, \underline{\Delta}_2), \dots$ , which are independent pairs of random vectors, each  $\underline{\Delta}_i$  is distributed as  $G$  on  $\Omega$  and  $\underline{x}_i = (X_{i1}, \dots, X_{ik})$  has conditional density function  $f(\underline{x}|\underline{\theta})$  given  $\underline{\Delta}_i = \underline{\theta}$ . The empirical Bayes approach, which was introduced by Robbins (1955), attempts to construct a decision rule concerning  $\underline{\Delta}_{n+1}$  at stage  $n+1$  based on  $\underline{x}_1, \dots, \underline{x}_{n+1}$ . The risk at stage  $n+1$  taking action  $\delta_n(\underline{x}; \underline{x}_1, \dots, \underline{x}_n) = \delta_n(\underline{x})$  is given by

$$\begin{aligned}
(3.9) \quad r_n(G, \delta_n) &= \int_{\mathcal{X}^n} E_n \left\{ \sum_{i \in \delta_n(\underline{x})} \left[ \int_{\Omega} (\theta_0 - \Delta - \theta_i) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) + \right. \right. \\
&\quad \left. \left. 2 \int_{(\theta_0, \infty)} (\theta_i - \theta_0) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) \right] \right\} d\underline{x} + r(G, \phi),
\end{aligned}$$

where  $E_n$  denotes the expectation with respect to the  $n$  independent random variables  $X_1, \dots, X_n$  each with common density function

$$m(\underline{x}) = \int_{\Omega} f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) = \prod_{i=1}^k m_i(x_i).$$

**Definition 3.1.** The sequence of procedures  $\{\delta_n\}$  is said to be asymptotically optimal (a.o.) relative to  $G$  if  $r_n(G, \delta_n) - r(G) = o(1)$  as  $n \rightarrow \infty$ , where  $r(G) = \inf_{\delta} r(G, \delta)$ .

In order to find an a.o. sequences of rules, let

$$\delta_{1,B}(\underline{x}) = \{i | x_i \leq \theta_0, \Delta_{1,G_i}(x_i) < 0\} \text{ and } \delta_{2,B}(\underline{x}) = \{i | \theta_0 < x_i < \theta_0 + \Delta, \Delta_{2,G_i}(x_i) < 0\}.$$

$$\delta_B(\underline{x}) = \delta_{1,B}(\underline{x}) \cup \delta_{2,B}(\underline{x}). \text{ For any } i = 1, 2, \dots, k \text{ and } \ell = 1, 2, \text{ let}$$

$\Delta_{\ell,i,n}(x_i) = \Delta_{\ell,i}(x_i; x_{1i}, \dots, x_{ni})$ ,  $n = 1, 2, \dots$  be two sequences of real-valued measurable functions. We define

$$(3.10) \quad \delta_n(\underline{x}) = \delta_{1,n}(\underline{x}) \cup \delta_{2,n}(\underline{x}),$$

where

$$\delta_{1,n}(\underline{x}) = \{i | x_i \leq \theta_0, \Delta_{1,i,n}(x_i) < 0\}$$

and

$$\delta_{2,n}(\underline{x}) = \{i | \theta_0 < x_i < \theta_0 + \Delta, \Delta_{2,i,n}(x_i) < 0\}.$$

We have the following theorem:

**Theorem 3.1.** If  $\int_0^{\infty} \theta dG_i(\theta) < \infty$ ,  $i = 1, 2, \dots, k$  and  $\Delta_{1,i,n}(x_i) \xrightarrow{P} \Delta_{1,G_i}(x_i)$ ,

for almost all  $x_i \leq \theta_0$  and  $\Delta_{2,i,n}(x_i) \xrightarrow{P} \Delta_{2,G_i}(x_i)$ , for almost all

$\theta_0 < x_i < \theta_0 + \Delta$ , where " $\xrightarrow{P}$ " means convergence in probability. Then  $\{\delta_n(\underline{x})\}$  defined by (3.10) is a.o. relative to  $G$ .

Proof.  $0 \leq \int_{\Omega} L(\underline{\theta}, \delta_n(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) - \int_{\Omega} L(\underline{\theta}, \delta_B(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta})$

$$(3.11) = \left\{ \sum_{i \in \delta_{1,n}(\underline{x})} \Delta_{1,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) - \sum_{i \in \delta_{1,B}(\underline{x})} \Delta_{1,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \right\} \\ + \left\{ \sum_{i \in \delta_{2,n}(\underline{x})} \Delta_{2,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) - \sum_{i \in \delta_{2,B}(\underline{x})} \Delta_{2,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \right\}.$$

The first term of (3.11) can be expressed as

$$(3.12) \left\{ \sum_{i \in \delta_{1,n}(\underline{x})} \Delta_{1,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) - \sum_{i \in \delta_{1,n}(\underline{x})} \Delta_{1,i,n}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \right\} \\ + \left\{ \sum_{i \in \delta_{1,n}(\underline{x})} \Delta_{1,i,n}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) - \sum_{i \in \delta_{1,B}(\underline{x})} \Delta_{1,i,n}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \right\} \\ + \left\{ \sum_{i \in \delta_{1,B}(\underline{x})} \Delta_{1,i,n}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) - \sum_{i \in \delta_{1,B}(\underline{x})} \Delta_{1,G_i}(\underline{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \right\} \\ \leq \sum_{i \in \delta_{1,n}(\underline{x})} (\Delta_{1,G_i}(\underline{x}_i) - \Delta_{1,i,n}(\underline{x}_i)) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j) \\ + \sum_{i \in \delta_{1,B}(\underline{x})} (\Delta_{1,i,n}(\underline{x}_i) - \Delta_{1,G_i}(\underline{x}_i)) \prod_{\substack{j=1 \\ j \neq i}}^k m_j(\underline{x}_j).$$

Since by the definition of  $\delta_{1,n}(\underline{x})$ , the second sum of (3.12) is less than or equal to zero.

The second term of (3.11) has a similar result.

Hence, if  $\Delta_{\ell,i,n}(\underline{x}_i) \xrightarrow{P} \Delta_{\ell,G_i}(\underline{x}_i)$ ,  $\ell = 1, 2$ , then



$$\begin{aligned}
0 &\leq \int_{\Omega} L(\underline{\theta}, \delta_n(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) - \int_{\Omega} L(\underline{\theta}, \delta_B(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) \\
&\leq 2 \sum_{i=1}^k |\Delta_{1,G_i}(x_i) - \Delta_{1,i,n}(x_i)| \prod_{\substack{j=1 \\ j \neq i}}^k m_j(x_j) + 2 \sum_{i=1}^k |\Delta_{2,G_i}(x_i) - \Delta_{2,i,n}(x_i)| \prod_{\substack{j=1 \\ j \neq i}}^k m_j(x_j) \\
&\leq 4\epsilon \sum_{i=1}^k \left( \prod_{\substack{j=1 \\ j \neq i}}^k m_j(x_j) \right)
\end{aligned}$$

with probability near 1, for large  $n$ . Hence

$$\int_{\Omega} L(\underline{\theta}, \delta_n(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta}) \xrightarrow{P} \int_{\Omega} L(\underline{\theta}, \delta_B(\underline{x})) f(\underline{x}|\underline{\theta}) dG(\underline{\theta})$$

for almost all  $\underline{x}$ .

By Corollary 1 of Robbins (1964),  $\{\delta_n(\underline{x})\}$  is a.o. relative to  $G$ .

From Theorem 3.1, our problem is reduced to finding consistent estimators of  $\Delta_{1,G_i}(x_i)$  and  $\Delta_{2,G_i}(x_i)$ . Let

$$(3.13) \quad M_{in}(x_i) = \frac{1}{n} \sum_{j=1}^n I_{(-\infty, x_i]}(X_{ji}),$$

then  $M_{in}(x_i) \xrightarrow{P} M_i(x_i)$  for all  $x_i > 0$ . Next, let  $\varphi(x) \geq 0$  be a Borel function satisfying the following conditions:

$$(3.14) \quad (i) \quad \sup_{-\infty < x < \infty} \varphi(x) < \infty, \quad (ii) \quad \int_{-\infty}^{\infty} \varphi(x) dx = 1, \quad \text{and} \quad (iii) \quad \lim_{x \rightarrow \infty} x\varphi(x) = 0$$

and  $\{h(n)\}$  be a sequence of positive constants satisfying the following conditions:

$$(3.15) \quad (i) \quad h(n) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ and } (ii) \quad nh(n) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

We define

$$(3.16) \quad m_{in}(x) = \frac{1}{nh(n)} \sum_{j=1}^n \varphi\left(\frac{x - X_{ji}}{h(n)}\right).$$

then  $m_{in}(x) \xrightarrow{P} m_i(x)$  for all  $x$  (see Parzen (1962)). For  $i = 1, 2, \dots, k$ , let

$$(3.17) \quad \Delta_{1,i,n}(x_i) = (\theta_0 - \Delta - x_i)m_{in}(x_i) + 1 - 2M_{in}(\theta_0) + M_{in}(x_i)$$

and

$$(3.18) \quad \Delta_{2,i,n}(x_i) = (x_i - \theta_0 - \Delta)m_{in}(x_i) + 1 - M_{in}(x_i).$$

Then

$$\Delta_{1,i,n}(x_i) \xrightarrow{P} \Delta_{1,G_i}(x_i) \quad \text{for all } x_i \leq \theta_0$$

and

$$\Delta_{2,i,n}(x_i) \xrightarrow{P} \Delta_{2,G_i}(x_i) \quad \text{for all } \theta_0 < x_i < \theta_0 + \Delta.$$

Finally, we define

$$\delta_n(\underline{x}) = \{i | x_i \leq \theta_0, \Delta_{1,i,n}(x_i) < 0\} \cup \{i | \theta_0 < x_i < \theta_0 + \Delta, \Delta_{2,i,n}(x_i) < 0\}.$$

Then  $\{\delta_n(\underline{x})\}$  is a.o. relative to  $G$ .

### 3.2. $\theta_0$ unknown

If  $\theta_0$  is unknown, let  $\pi_0$  be the control population and  $X_0$  be the random variable from  $\pi_0$ . We assume that  $X_0$  has conditional pdf

$$f(x_0 | \theta_0) = \frac{1}{\theta_0} I_{(0, \theta_0)}(x_0), \quad \theta_0 > 0. \quad \text{In this case}$$

$$\Omega = \{\underline{\theta} = (\theta_0, \theta_1, \dots, \theta_k) | \theta_i > 0, i = 0, 1, \dots, k\}, \quad \mathcal{X} = \{\underline{x} = (x_0, x_1, \dots, x_k)$$

$$| x_i > 0, i = 0, 1, \dots, k\}, \quad G(\underline{\theta}) = \prod_{i=0}^k G_i(\theta_i), \quad f(\underline{x} | \underline{\theta}) = \prod_{i=0}^k f(x_i | \theta_i), \quad \text{and}$$

at stage  $n$  we observed  $\underline{x}_n = (x_{n0}, x_{n1}, \dots, x_{nk})$ . Under the loss function

(2.4), the Bayes rule  $\delta_B(\underline{x})$  is given by

$$i \in \delta_B(\underline{x}) \text{ if } \Delta_{G_0, G_i}(x_0, x_i) < 0$$

where

$$\Delta_{G_0, G_i}(x_0, x_i) = \int \theta_0 f(x_0 | \theta_0) dG_0(\theta_0) m_i(x_i) - \Delta m_0(x_0) m_i(x_i) -$$

$$\int \theta_i f(x_i | \theta_i) dG_i(\theta_i) m_0(x_0) + 2 \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) f(x_i | \theta_i) f(x_0 | \theta_0) dG_i(\theta_i) dG_0(\theta_0).$$

Using formula (3.4) if  $0 < x_i \leq x_0$ , we have

$$\begin{aligned} & \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) f(x_i | \theta_i) f(x_0 | \theta_0) dG_i(\theta_i) dG_0(\theta_0) \\ &= m_0(x_0) - \int_{x_0}^{\infty} \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0), \end{aligned}$$

and if  $0 < x_0 < x_i$ , we have

$$\begin{aligned} & \int_{\{\theta_0 < \theta_i\}} (\theta_i - \theta_0) f(x_i | \theta_i) f(x_0 | \theta_0) dG_i(\theta_i) dG_0(\theta_0) \\ &= (1 - G_i(x_i))(m_0(x_0) - m_0(x_i)) - m_i(x_i)(G_0(x_i) - G_0(x_0)) \\ & \quad - \int_{x_i}^{\infty} \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) + m_0(x_i). \end{aligned}$$

Hence

$$\begin{aligned} (3.19) \quad \Delta_{G_0, G_i}(x_0, x_i) &= m_i(x_i)(1 - M_0(x_0)) + (1 + M_i(x_i))m_0(x_0) + \\ & \quad (x_0 - x_i - \Delta)m_i(x_i)m_0(x_0) - 2 \int_{x_0}^{\infty} \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \\ &= \Delta_{1, G_0, G_i}(x_0, x_i) \text{ (say), if } 0 < x_i \leq x_0 \end{aligned}$$

and

$$\begin{aligned} (3.20) \quad \Delta_{G_0, G_i}(x_0, x_i) &= (1 - M_i(x_i))m_0(x_0) + (1 + M_0(x_0) - 2M_0(x_i))m_i(x_i) + \\ & \quad (x_i - x_0 - \Delta)m_i(x_i)m_0(x_0) + 2M_i(x_i)m_0(x_i) - 2 \int_{x_i}^{\infty} \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \\ &= \Delta_{2, G_0, G_i}(x_0, x_i) \text{ (say), if } 0 < x_0 < x_i. \end{aligned}$$

Thus

$$(3.21) \quad \delta_B(\underline{x}) = \delta_{1, B}(\underline{x}) \cup \delta_{2, B}(\underline{x})$$

where

$$\delta_{1,B}(\underline{x}) = \{i | 0 < x_i \leq x_0, \Delta_{1,G_0,G_i}(x_0, x_i) < 0\}$$

and

$$\delta_{2,B}(\underline{x}) = \{i | 0 < x_0 < x_i, \Delta_{2,G_0,G_i}(x_0, x_i) < 0\}.$$

Similar to Theorem 3.1, we have the following result.

Theorem 3.2. If  $\int_0^\infty \theta dG_i(\theta) < \infty$ ,  $i = 0, 1, \dots, k$  and for all  $1 \leq i \leq k$ ,

$$\Delta_{1,i,n}(x_0, x_i) \xrightarrow{P} \Delta_{1,G_0,G_i}(x_0, x_i) \text{ for } x_i \leq x_0 \text{ and}$$

$$\Delta_{2,i,n}(x_0, x_i) \xrightarrow{P} \Delta_{2,G_0,G_i}(x_0, x_i) \text{ for } x_0 < x_i. \text{ Let}$$

$$\delta_n(\underline{x}) = \{i | x_i \leq x_0, \Delta_{1,i,n}(x_0, x_i) < 0\} \cup \{i | x_0 < x_i, \Delta_{2,i,n}(x_0, x_i) < 0\},$$

then  $\{\delta_n(\underline{x})\}$  is a.o. relative to  $G$ .

Hence our problem is to find a consistent estimator of

$$\int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \text{ for } x_0 \leq a.$$

Theorem 3.3. Let  $M_{in}(x)$  and  $m_{in}(x)$  be defined by (3.13) and (3.16), respectively. Then

$$\int_a^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0) \xrightarrow{P} \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \text{ for } x_0 \leq a,$$

where  $G_{0n}(\theta_0) = M_{0n}(\theta_0) - \theta_0 m_{0n}(\theta_0)$ .

$$\begin{aligned} \text{Proof. } & \left| \int_a^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0) - \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_{0n}(\theta_0) \right| \\ & \leq \int_a^\infty \frac{|M_{in}(\theta_0) - M_i(\theta_0)|}{\theta_0} dG_{0n}(\theta_0) \\ & \leq \frac{1}{a} \sup_{-\infty < x < \infty} |M_{in}(x) - M_i(x)| \leq \epsilon \end{aligned}$$

with probability near 1, for large  $n$ , by Glivenko-Cantelli Theorem. Since

$\frac{M_i(\theta_0)}{\theta_0}$  is bounded continuous and  $G_{0n}(\theta_0) \xrightarrow{P} G_0(\theta_0)$ , we have

$$\int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_{0n}(\theta_0) \xrightarrow{P} \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0).$$

Thus

$$\begin{aligned} & \left| \int_a^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0) - \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \right| \\ & \leq \left| \int_a^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0) - \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_{0n}(\theta_0) \right| + \\ & \quad \left| \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_{0n}(\theta_0) - \int_a^\infty \frac{M_i(\theta_0)}{\theta_0} dG_0(\theta_0) \right| \\ & \leq \varepsilon \text{ with probability near 1, for large } n. \end{aligned}$$

From Theorem 3.3, if we define

$$\begin{aligned} \Delta_{1,i,n}(x_0, x_i) &= m_{in}(x_i)(1 - M_{0n}(x_0)) + m_{0n}(x_0)(1 + M_{in}(x_i)) \\ &+ (x_0 - x_i - \Delta)m_{in}(x_i) - m_{0n}(x_0) - 2 \int_{x_0}^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0), \end{aligned}$$

where  $G_{0n}(\theta_0) = M_{0n}(\theta_0) - \theta_0 m_{0n}(\theta_0)$  and  $M_{in}(x)$ ,  $m_{in}(x)$  are defined by (3.13) and (3.16), respectively, and

$$\begin{aligned} \Delta_{2,i,n}(x_0, x_i) &= m_{0n}(x_0)(1 - M_{in}(x_i)) + m_{in}(x_i)(1 + M_{0n}(x_0) - 2M_{0n}(x_i)) \\ &+ (x_i - x_0 - \Delta)m_{0n}(x_0)m_{in}(x_i) + 2M_{in}(x_i)m_{0n}(x_i) - 2 \int_{x_i}^\infty \frac{M_{in}(\theta_0)}{\theta_0} dG_{0n}(\theta_0). \end{aligned}$$

Then

$$\Delta_{\ell,i,n}(x_0, x_i) \xrightarrow{P} \Delta_{\ell,G_0,G_i}(x_0, x_i), \quad \ell = 1, 2.$$

Now, let

$$\delta_n(\underline{x}) = \{i | x_i \leq x_0, \Delta_{1,i,n}(x_0, x_i) < 0\} \cup \{i | x_0 < x_i, \Delta_{2,i,n}(x_0, x_i) < 0\}.$$

From Theorem 3.2, we have  $\{\delta_n(\underline{x})\}$  is a.o. relative to  $G$ .

### 3.3. Rate of Convergence of the Empirical Bayes Rules

In this section we will consider the rate of convergence of the empirical Bayes rules derived in Section 3.1.

**Definition 3.2.** The sequence of procedures  $\{\delta_n\}$  is said to be asymptotically optimal of order  $\alpha_n$  relative to  $G$  if  $r_n(G, \delta_n) - r(G) = O(\alpha_n)$  as  $n \rightarrow \infty$ , where  $\lim_{n \rightarrow \infty} \alpha_n = 0$ .

The main result (Theorem 3.8) of this section is based on a series of lemmas.

**Lemma 3.4.** Let  $\Delta_{1,G_i}(x_i)$ ,  $\Delta_{2,G_i}(x_i)$ ,  $\Delta_{1,i,n}(x_i)$  and  $\Delta_{2,i,n}(x_i)$  be defined by (3.6), (3.7), (3.17) and (3.18) respectively. Then  $0 \leq r_n(G, \delta_n) - r(G)$

$$\leq \sum_{i=1}^k \int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} E|\Delta_{1,i,n}(x_i) - \Delta_{1,G_i}(x_i)|^\delta dx_i + \\ \sum_{i=1}^k \int_{\theta_0}^{\theta_0+\Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} E|\Delta_{2,i,n}(x_i) - \Delta_{2,G_i}(x_i)|^\delta dx_i, \quad \delta > 0.$$

**Proof.** The proof is similar to that of Lemma 3 of Van Ryzin and Susarla (1977) and hence omitted.

**Lemma 3.5.** Let  $\varphi(x)$  satisfy the conditions (i)  $\varphi(x) = 0$  if  $x \notin (0, a)$  for some finite  $a > 0$ , (ii)  $\int_0^a \varphi(x) dx = 1$ , and (iii)  $\sup_x |\varphi(x)| < \infty$  and define

$m_{in}(x_i) = \frac{1}{nh(n)} \sum_{j=1}^n \varphi\left(\frac{X_{ji} - x_i}{h(n)}\right)$ , where  $\{h(n)\}$  satisfy the conditions (3.15) (see Johns and Van Ryzin (1972)). Then

$|Em_{in}(x_i) - m_i(x_i)| \leq h(n) f_\epsilon(x_i) \int_0^a |u \varphi(u)| du$ , for large  $n$ , where

$f_\epsilon(x_i) = \sup_{0 \leq y \leq \epsilon} |m'_i(x_i + y)|$ ,  $\epsilon > 0$ .

Proof.  $E m_{in}(x_i) - m_i(x_i)$

$$\begin{aligned}
 &= \frac{1}{h(n)} \int \varphi\left(\frac{y-x_i}{h(n)}\right) m_i(y) dy - m_i(x_i) \\
 &= \int_0^a \varphi(u) [m_i(x_i + uh(n)) - m_i(x_i)] du \\
 &= \int_0^a \varphi(u) [uh(n) m'_i(x_i + \eta_n(x_i, u))] du
 \end{aligned}$$

where  $0 < \eta_n(x_i, u) < uh(n)$ .

For  $\epsilon > 0$ , let  $n$  be large enough so that  $uh(n) \leq \epsilon$ , then

$$|E m_{in}(x_i) - m_i(x_i)| \leq h(n) f'_\epsilon(x_i) \int_0^a |u \varphi(u)| du.$$

Lemma 3.6. Under the conditions of Lemma 3.5, we have

$$\text{var } m_{in}(x_i) \leq \frac{1}{nh(n)} m_i(x_i) \int_0^a \varphi^2(u) du.$$

Proof.  $\text{Var } m_{in}(x_i) = \text{var} \left\{ \frac{1}{nh(n)} \sum_{j=1}^n \varphi\left(\frac{x_{ji} - x_i}{h(n)}\right) \right\}$

$$\leq \frac{1}{nh(n)} \int_0^a \varphi^2(u) m_i(x_i + uh(n)) du$$

$$\leq \frac{1}{nh(n)} m_i(x_i) \int_0^a \varphi^2(u) du, \text{ since } m_i(x_i) +.$$

Remark: From Lemma 3.5 and Lemma 3.6, we have

$$m_{in}(x_i) \xrightarrow{P} m_i(x_i) \text{ if } f'_\epsilon(x_i) < \infty.$$

Lemma 3.7. Under the conditions of Lemma 3.5, we have

$$(a) \text{ Var } \Delta_{1,i,n}(x_i) = O((\theta_0 - \Delta - x_i)^2 m_i(x_i) \frac{1}{nh(n)}),$$

$$(b) \text{ Var } \Delta_{2,i,n}(x_i) = O((x_i - \theta_0 - \Delta)^2 m_i(x_i) \frac{1}{nh(n)}).$$

Proof. (a)  $\text{Var } \Delta_{1,i,n}(x_i)$

$$\begin{aligned} &\leq 2\{(\theta_0 - \Delta - x_i)^2 \text{Var } m_{in}(x_i) + \text{var}(M_{in}(x_i) - 2M_{in}(\theta_0))\} \\ &\leq 2\{(\theta_0 - \Delta - x_i)^2 m_i(x_i) \frac{1}{nh(n)} \int_0^a \varphi^2(u) du + \frac{5}{2n}\} \text{ (By Lemma 3.6)} \\ &\leq M \frac{1}{nh(n)} (\theta_0 - \Delta - x_i)^2 m_i(x_i), \text{ for some } M > 0. \end{aligned}$$

Similarly, we have the result (b).

**Theorem 3.8.** Under the conditions of Lemma 3.5. If

$$(i) \int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} |\theta_0 - \Delta - x_i|^{\delta} m_i^{\delta/2}(x_i) dx_i < \infty,$$

$$(ii) \int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} |x_i - \theta_0 - \Delta|^{\delta} m_i^{\delta/2}(x_i) dx_i < \infty,$$

$$(iii) \int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} |\theta_0 - \Delta - x_i|^{\delta} f_{\epsilon}^{\delta}(x_i) dx_i < \infty,$$

and

$$(iv) \int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} |x_i - \theta_0 - \Delta|^{\delta} f_{\epsilon}^{\delta}(x_i) dx_i < \infty,$$

where  $0 < \delta < 2$ , then

$$r_n(G, \delta_n) - r(G) = O(\max\{(\frac{1}{nh(n)})^{\delta/2}, (h(n))^{\delta}\}) \text{ as } n \rightarrow \infty.$$

Proof. For  $0 < \delta < 2$ , by Hölder inequality and Lemma 3.4, we have

$$\begin{aligned} 0 &\leq r_n(G, \delta_n) - r(G) \\ &\leq \sum_{i=1}^k \{\max(1, 2^{\delta-1}) [\int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} (\text{Var } \Delta_{1,i,n}(x_i))^{\delta/2} dx_i + \end{aligned}$$



$$\begin{aligned}
& \int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} |(\theta_0 - \Delta - x_i)(E m_{in}(x_i) - m_i(x_i))|^\delta dx_i \} + \\
& \sum_{i=1}^k \{ \max(1, 2^{\delta-1}) [\int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} (\text{var } \Delta_{2,i,n}(x_i))^{\delta/2} dx_i + \\
& \int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} |(x_i - \theta_0 - \Delta)(E m_{in}(x_i) - m_i(x_i))|^\delta dx_i] \}.
\end{aligned}$$

By Lemma 3.7, we have

$$\int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} (\text{var } \Delta_{1,i,n}(x_i))^{\delta/2} dx_i = O((nh(n))^{-\delta/2})$$

and

$$\int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} (\text{var } \Delta_{2,i,n}(x_i))^{\delta/2} dx_i = O((nh(n))^{-\delta/2}).$$

By Lemma 3.5, we have

$$\int_0^{\theta_0} |\Delta_{1,G_i}(x_i)|^{1-\delta} |\theta_0 - \Delta - x_i|^\delta |E m_{in}(x_i) - m_i(x_i)|^\delta dx_i = O((h(n))^\delta)$$

and

$$\int_{\theta_0}^{\theta_0 + \Delta} |\Delta_{2,G_i}(x_i)|^{1-\delta} |(x_i - \theta_0 - \Delta)(E m_{in}(x_i) - m_i(x_i))|^\delta dx_i = O((h(n))^\delta).$$

Hence

$$r_n(G, \delta_n) - r(G) = O(\max\{(nh(n))^{-\delta/2}, (h(n))^\delta\}) \text{ as } n \rightarrow \infty.$$

**Corollary 3.9.** Under the conditions of Theorem 3.8. If we take  $h(n) = n^{-\alpha}$ ,  $0 < \alpha < 1$ , then the optimal choice of  $\alpha$  is  $1/3$  and  $r_n(G, \delta_n) - r(G) = O(n^{-\delta/3})$  as  $n \rightarrow \infty$ .

Remark: If the prior distribution  $G_j$  is such that  $g_j(x)/x$  and  $m_j(x)$  are both bounded on  $(0, \theta_0 + \Delta + \epsilon)$ , it is easy to check that the conditions of Theorem 3.8 are satisfied for  $0 < \delta \leq 1$ .

## REFERENCES

1. Deely, J. J. (1965). Multiple decision procedures from an empirical Bayes approach. Ph.D. Thesis (Mimeo. Ser. No. 45), Dept. of Statist., Purdue University.
2. Gupta, S. S. and Hsiao, P. (1981). Empirical Bayes rules for selecting good populations. Mimeo. Ser. No. 81-5, Dept. of Statist., Purdue University.
3. Johns, M. V. Jr. and Van Ryzin, J. (1972). Convergence rates for empirical Bayes two-action problems II, continuous case. Ann. Math. Statist. 43, 934-947.
4. Parzen, E. (1962). On estimation of a probability density function and mode. Ann. Math. Statist. 33, 1065-1076.
5. Robbins, H. (1955). An empirical Bayes approach to statistics. Proc. 3rd Berkeley Symp. Math. Statist. Prob. 155-163.
6. Robbins, H. (1964). The empirical Bayes approach to statistical decision problems. Ann. Math. Statist. 35, 1-20.
7. Singh, A. K. (1977). On slippage tests and multiple decision (selection and ranking) procedures. Ph.D. Thesis (Mimeo. Ser. No. 494), Dept. of Statist., Purdue University.
8. Van Ryzin, J. (1970). On some nonparametric empirical Bayes multiple decision problems. Proc. First Internat. Symp. Nonparametric Techniques in Statistic Inference (Ed. M. L. Puri), 585-603.
9. Van Ryzin, J. and Susarla, V. (1977). On the empirical Bayes approach to multiple decision problems. Ann. Statist. 5, 172-181.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report #83-37	2. GOVT ACCESSION NO. AD-A133336	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON BAYES AND EMPIRICAL BAYES RULES FOR SELECTING GOOD POPULATIONS		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Shanti S. Gupta and Lii-Yuh Leu		6. PERFORMING ORG. REPORT NUMBER Technical Report #83-37
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, IN 47907		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0455
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1983
		13. NUMBER OF PAGES 21
		15. SECURITY CLASS. (of this report) Unclassified
		16. DECLASSIFICATION, DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bayes rules, empirical Bayes rules, selection procedures, asymptotically optimal, rate of convergence.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper deals with the problem of selecting all populations which are close to a control or standard. A general Bayes rule for the above problem is derived. Empirical Bayes rules are derived when the populations are assumed to be uniformly distributed. Under some conditions on the marginal and prior distributions, the rate of convergence of the empirical Bayes risk to the minimum Bayes risk is investigated. The rate of convergence is shown to be $n^{-\delta/3}$ for some $\delta$ , $0 < \delta < 2$ .		

EN

DA

FIL

10

DT